arXiv:1211.1983v2 [hep-th] 21 Dec 2012

QMUL-PH-12-15

The Navier-Stokes equation and solution generating symmetries from holography

Joel Berkeley[1] and David S. Berman[2],

Queen Mary University of London, Centre for Research in String Theory,
School of Physics, Mile End Road, London, E1 4NS, England

Abstract

The fluid-gravity correspondence provides us with explicit spacetime metrics that are holographically dual to (non-)relativistic nonlinear hydrodynamics. The vacuum Einstein equations, in the presence of a Killing vector, possess solution-generating symmetries known as Ehler's transformations. We apply these Ehler's transformations to vacuum metrics with hydrodynamic duals to develop a formalism for solution-generating transformations of Navier-Stokes fluids. Using this we provide examples of a linear energy scaling from RG flow under vanishing vorticity, and a set of $\mathbb{Z}_2$ symmetries for fixed viscosity.

---

[1] `j.a.fitzhardinge-berkeley@qmul.ac.uk`
[2] `d.s.berman@qmul.ac.uk`

Contents

1 Introduction **1**

2 Hydrodynamics **3**
  2.1 Relativistic fluids 4
  2.2 Non-relativistic Navier-Stokes fluids 6
  2.3 The Navier-Stokes fluid on a Rindler boundary 7
  2.4 The Navier-Stokes fluid on a Schwarzschild sphere 11

3 Duality in the context of holography **12**
  3.1 Symmetry groups of the Einstein equations 13
  3.2 The spacetime Ehlers group 14

4 Solution-generating the Navier-Stokes fluid **15**
  4.1 Transforming the fluid 16
    4.1.1 Energy scaling invariance from a bulk isometry 17
    4.1.2 The timelike Killing vector 17
  4.2 Fixed viscosity dualities 17
    4.2.1 Spacelike Killing vectors 18
    4.2.2 Group structure for fixed boundary 19

5 Discussion **20**

1 Introduction

In 1974, Damour [1], and later in 1986, Thorne et al. [2], considered an observer outside a black hole, interacting with (perturbing) the event horizon. Surprisingly, they found that the observer will experience the perturbations of the "stretched" horizon as modes of a viscous fluid possessing electric charge and conductivity. This inspired a host of works over the years but the connection between gravitational physics and fluids became sharper with the advent of the AdS/CFT correspondence when Policastro et. al [5] related the shear viscosity of $\mathcal{N} = 4$ super Yang-Mills theory to the absorption of energy by a black brane.

This was the start of using the holographic principle, that is the correspondence between gravitational theories on $(d+1)$-dimensional manifolds and d-dimensional quantum field theories, as a tool for calculating hydrodynamic properties.

More recently, there has been a set of works that directly relate solutions of Einstein's equations of a particular type to solutions to the Navier-Stokes equations in one dimension less [6, 7, 8, 9, 10, 11, 12, 13, 14]. Later we will review the details of how this correspondence is derived but the essential flavour is as follows. One writes down a very particular ansatz for the metric in $d+1$ dimensions which has undetermined functions, $v^i(x,t), P(x,t)$ and parameter, ν with the index $i = 1, .., d-1$ i.e. over a $(d-1)$ subset of the d spacetime dimensions. Solving the Einstein equations then constrains the functions $v^i(x,t), P(x,t)$ to give a set of second order nonlinear differential equations for $v^i(x,t), P(x,t)$. This set of equations are the Navier-Stokes equations describing a fluid in d dimensions with pressure, $P(x,t)$, fluid velocity field $v^i(x,t)$ and viscosity ν. Thus particular solutions to the Navier-Stokes equations provide particular solutions to the Einstein equations.

Other recent work on the fluid/gravity correspondence may be found in [20, 21, 22, 23, 24, 25, 26, 27, 28].

It has been known since Buchdahl [17] that for manifolds with isometries Einstein's equations have solution generating symmetries. That is there are "hidden" symmetries of the equations that map from one solution to the other. In fact there is a vast set of these as described by Ehler [18] and Geroch [19]. The question we wish to pose in this paper is whether the solution generating symmetries of Einstein's equations can lead to solution generating symmetries in the Navier-Stokes equations?

The procedure to determine this will be as follows:

- impose a Killing symmetry in a spacetime that admits the metric ansatz corresponding to the Navier-Stokes equations;
- carry out Ehler's transformations that preserve the ansatz;
- determine the induced transformation of the Navier-Stokes data i.e. v^i, P, ν.

The procedure could immediately fail if there were no Ehler's transformations that preserve the metric ansatz required for the fluid/gravity correspondence. We will find that there are a finite set and we will be able to explore the transformations on the Navier-Stokes fields for different choices of Killing directions in spacetime.

The paper will try to be as self contained as possible and so we begin with a review of the necessary ideas in fluids; the Navier-Stokes equation; the fluid gravity correspondence; and solution generating symmetries in general relativity. We will then carry out the procedure described above for spatial, timelike and null Killing vectors to see what solution generating symmetries they correspond to in the Navier-Stokes equation. We end with some comments and ideas for future work. A reader familiar with the formalism of hydrodynamics and the Navier-Stokes equation may wish to skip directly to section 3 where we carry out the duality transformation in the gravity dual to see the induced transformations.

2 Hydrodynamics

The study of hydrodynamics is fundamental to vast areas of physics and engineering, owing to its origin as the long-wavelength limit of any interacting field theory at finite temperature. Such a limit needs a consistent definition. Consider a quantum field theory where quanta interact with a characteristic length scale ℓ_{corr}, the correlation length. The long-wavelength limit simply requires that fluctuations of the thermodynamic quantities of the system vary with a length scale L much greater than ℓ_{corr}, parameterized by the dimensionless Knudsen number

$$K_n = \frac{\ell_{\text{corr}}}{L}. \tag{1}$$

For a physical description to be useful in non-equilibrium states, we naturally require that L remain small compared to the size of the system. This is usually satisfied trivially by considering systems of infinite size.

The long-wavelength limit allows the definition of a particle as an element of the macroscopic fluid, infinitesimal with respect to the size of the system, yet containing a sufficiently large number of microscopic quanta. One mole contains an Avogadro's number of molecules, for example. Each particle defines a local patch of the fluid in thermal equilibrium, that is, thermodynamic quantities do not vary within the particle. Away from global equilibrium quantities vary between particles as function of time τ and spatial coordinates $\vec{x}$, combined as $x^a = (\tau, \vec{x})$. The evolution of the quantities between particles is parameterized by a relativistic velocity $u^b(x^a)$, which refers to the velocity of the fluid at x^a, rather than the velocity of an individual particle as it traverses a path through the fluid, parameterized by the coordinates. It is well known [29] that the thermodynamic quantities, such as the temperature $T(x^a)$ and the density $\rho(x^a)$, are determined by the value of any two of them, along with the equation of state. The evolution of the system is then specified by the equations of hydrodynamics in terms of a set of transport coefficients, whose values depend on the fluid in question.

Fluid flow is in general relativistic in that the systems it describes are constrained by local Lorentz invariance, and velocities may take any physical values below the speed of light. Applications at relativistic velocities are multitudinous: the dust clouds in galaxy and star formation; the flow of plasmas and gases in stars supporting fusion; the superfluid cores of neutron stars; the horizons of black holes are all described by hydrodynamics. Modelling black holes (and black branes in M/string theory) with hydrodynamics has now developed into a fundamental correspondence of central importance to our present study, we elaborate in §1. Quark-gluon plasmas behave as nearly ideal fluids and are expected to have formed after the inflationary epoch of the big bang, are reproduced in collisions at the RHIC and LHC, and in the core of nuclear fusion reactors. Non-relativistic fluids are equally ubiquitous, somewhat more familiar, and constitute an endless list of phenomena from the atmosphere to the oceans.

2.1 Relativistic fluids

We begin with a discussion, adapted from [30], of the relativistic fluid described by the stress energy tensor T^{ab} and a set of conserved currents J_I^a where I indexes the corresponding conserved charge. The dynamical equations of the d-spacetime dimensional fluid are

$$\nabla_a T^{ab} = 0 \qquad \nabla_a J_I^a = 0\,. \tag{2}$$

For an ideal fluid, with no dissipation, the energy-momentum tensor and currents may be expressed in a local rest frame in the form

$$T^{ab} = \rho u^a u^b + p(g^{ab} + u^a u^b) \tag{3a}$$

$$J_I^a = q_I u^a \tag{3b}$$

where p is the pressure and q_I the conserved charges. The velocity is normalised to $u^a u_a = -1$. The entropy current is given by (3b) with the charge q being given by the local entropy density. The conservation of the entropy current illustrates the non-dissipative nature intrinsic to zero entropy production.

In a dissipative fluid, there are corrections to (3). First, we must first take into account the interrelation between mass and energy to define the velocity field more rigorously. This is achieved by using the Landau gauge, which requires that the velocity be an eigenvector of the stress-energy tensor with eigenvalue the local energy density of the fluid (this is satisfied by the velocity normalisation for the ideal fluid). If the stress energy tensor gains a dissipative term Π^{ab}, and the current a term Υ^a, this reads

$$\Pi^{ab} u_a = 0 \qquad \Upsilon^a u_a = 0.$$

Dissipative corrections to the stress tensor are constructed in a derivative expansion of the velocity field and thermodynamic variables, where derivatives implicitly scale with the infinitesimal Knudsen number (1). Recalling that the equations of motion for the ideal fluid are composed of relations between these gradients, we may express Π^{ab} purely in terms of the derivative of the velocity. This can be iterated to all orders in the expansion. Now, the derivative of the velocity may be decomposed using the acceleration A^a, divergence θ, a symmetric traceless shear σ^{ab}, and the antisymmetric vorticity ω^{ab} into the form

$$\nabla^a u^b = -A^a u^b + \sigma^{ab} + \omega^{ab} + \frac{1}{d-1}\theta P^{ab},$$

where

$$\begin{aligned}\theta &= \nabla_a u^a \\ A^a &= u^b \nabla_b u^a \\ \sigma^{ab} &= P^{ac}P^{bd}\nabla_{(c}u_{d)} - \frac{1}{d-1}\theta P^{ab} \\ \omega^{ab} &= P^{bc}P^{ad}\nabla_{[c}u_{d]}.\end{aligned}$$

and $P^{ab} = g^{ab} + u^a u^b$ is a projection operator onto spatial directions. In the Landau frame, only the divergence and shear can be built purely from velocity gradients, and so define the leading order contribution to the stress-energy tensor. A similar analysis for the charge current retains the acceleration, and if one includes the parity-violating pseudo-vector contribution

$$\ell^a = \epsilon_{bcd}{}^a u^b \nabla^c u^d,$$

the leading order dissipative equations of motion for a relativistic fluid are (2) with

$$T^{ab} = \rho u^a u^b + pP^{ab} - 2\eta\sigma^{ab} - \zeta\theta P^{ab} \tag{4a}$$

$$J_I^a = q_I u^a - \chi_{IJ}P^{ab}\nabla_b q_J - \Theta_I \ell^a - \gamma_I P^{ab}\nabla_b T, \tag{4b}$$

where η and ζ are the shear[1] and bulk viscosities respectively, ζ_{IJ} is the matrix of charge diffusion coefficients, γ_I indicates the contribution of the energy density, and Θ_I the pseudo-vector transport coefficients. The transport coefficients have been calculated in the weakly coupled QFT in perturbation theory, whereas in the strongly coupled theory, a dual holographic description may be employed, see e.g. [3].

Returning to the discussion of the entropy current, we now expect that the entropy will increase in fluid evolution to equilibrium, consistent with the second law of thermodynamics. This translates to the requirement that

$$\nabla_a J_s^a = \frac{2\eta}{T}\sigma_{ab}\sigma^{ab} \geq 0,$$

which is satisfied iff the shear viscosity is positive, a condition satisfied for a large class of QFTs [4].

---

[1] We use the traditional notation of [4, 12, 29] rather than that of [8, 9]. We define the kinematical viscosity ν in §2.2.

2.2 Non-relativistic Navier-Stokes fluids

We now take the non-relativistic limit, defined by long distances, long times and low velocity and pressure amplitudes, of the relativistic fluid on a spacetime lightly perturbed in the following manner [12]. Consider an arbitrary perturbation

$$\hat{G}_{ab} = \hat{g}_{ab} + H_{ab}$$

of the background metric

$$\hat{g}_{ab} = -\mathrm{d}\tau^2 + \hat{g}_{ij}\mathrm{d}x^i\mathrm{d}x^j$$

(this form is purely a choice of coordinate system), where

$$\begin{aligned} H_{\tau\tau} &= \epsilon^2 h_{\tau\tau}(\epsilon^2\tau, \epsilon x^i) \\ H_{\tau i} &= \epsilon \hat{a}_i(\epsilon^2\tau, \epsilon x^i) \\ H_{ij} &= \epsilon^2 h_{ij}(\epsilon^2\tau, \epsilon x^i). \end{aligned} \tag{5}$$

Treating the fluid on $\hat{G}_{ab}$ as an effective forced flow of that on $\hat{g}_{ab}$, whilst preserving the normalisation $u_a u^a = -1$, the velocity

$$u^a = \frac{1}{\sqrt{1 - \hat{g}_{ij}V^iV^j}}(1, V^i),$$

density and pressure may be expanded about an equilibrium configuration (ρ_0, V_0^i, p_0) of a stationary, uniform fluid at rest, as

$$\begin{aligned} V^i &= \epsilon v^i(\epsilon^2\tau, \epsilon x^i) \\ \frac{p_e - p_0}{\rho_0 + p_0} &= \epsilon^2 p(\epsilon^2\tau, \epsilon x^i) \end{aligned} \tag{6}$$

with an infinitesimal scaling parameter ϵ. The scaling of the density is fixed by the equation of state. The scaling of time τ and space x^i are implicit. In this limit, the temporal component of the relativistic Navier-Stokes equations becomes

$$\nabla_a T^{a\tau} = \epsilon^2(\rho_0 + p_0)\nabla_i v^i + \mathcal{O}(\epsilon^4) \tag{7}$$

where ∇_i is the covariant derivative with respect to the spatial metric $\hat{g}_{ij}$. Separating the gauge field $\hat{a}_i$ into its pure curl and divergence parts

$$\hat{a}_i = a_i + \nabla_i\chi,$$

such that $\nabla_i a^i = 0$, and defining an effective pressure

$$P = p_e - \frac{1}{2}h_{\tau\tau} + \frac{\partial\chi}{\partial\tau},$$

the spatial components yield

$$\begin{aligned}\nabla_a T^{ai} =& \epsilon^3(\rho_0+p_0)[\nabla_i P + \partial_\tau v_i + v^j\nabla_j v_i - \nu(\nabla^j\nabla_j v_i + \hat{R}_{ij}v^j) \\ &+ \partial_\tau a_i + v^j f_{ji}] + \mathcal{O}(\epsilon^5)\end{aligned} \tag{8}$$

where

$$\nu = \eta/(\rho_0+p_0)$$

is the kinematical viscosity, $f_{ij} \equiv \partial_i a_j - \partial_j a_i$ is the field strength of a_i (equivalently $\hat{a}_i$) and $\hat{R}_{ij}$ is the Ricci tensor on $\hat{g}_{ij}$. If a fluid carries conserved charges, the equations $\nabla_a J^a_I = 0$ will also yield incompressibility (7). Taking the hydrodynamic limit $\epsilon \to 0$, these give the incompressible non-relativistic Navier-Stokes equations with a forcing function due to an external electromagnetic field,

$$\partial_\tau v_i - \nu(\partial^2 v_i + \hat{R}_{ij}v^j) + \partial_i P + v^j\partial_j v_i = -\partial_\tau a_i - v^j f_{ji} \tag{9a}$$

$$\partial_i v^i = 0. \tag{9b}$$

Ideal fluids are described by Euler's equations, obtained by setting $\nu = 0$. We will mostly be concerned with fluid flow on flat space ($\hat{R}_{ij} = 0$) in the absence of external forces, where H_{ab}, thus a_i, are zero.

2.3 The Navier-Stokes fluid on a Rindler boundary

A metric dual to the non-relativistic incompressible Navier-Stokes equations was first developed in [8] on the Rindler wedge. This included terms up to third order in the ϵ expansion as in (6). An algorithm for generalising this metric to all orders was subsequently developed in [9], though terms calculated beyond third order are not universal. They receive corrections from quadratic curvature in Gauss-Bonnet gravity [11]. We summarise the construction in [9] here.

Consider the surface Σ_c with induced metric

$$\gamma_{ab}\mathrm{d}x^a\mathrm{d}x^b = -r_c\mathrm{d}\tau^2 + \mathrm{d}x_i\mathrm{d}x^i \tag{10}$$

where the parameter $\sqrt{r_c}$ is an arbitrary constant. One metric embedding this surface is

$$\mathrm{d}s^2 = -r\mathrm{d}\tau^2 + 2\mathrm{d}\tau\mathrm{d}r + \mathrm{d}x_i\mathrm{d}x^i, \tag{11}$$

which describes flat space (fig. 1) in ingoing Rindler coordinates $x^\mu = (\tau, x^i, r)$, defined in terms of the Cartesian chart (t, x^i, z) by

$$z^2 - t^2 = 4r \qquad z + t = e^{\tau/2}. \tag{12}$$

The hypersurface Σ_c is defined by $r = r_c$ where r is the coordinate into the bulk. Allowing for a family of equilibrium configurations, consider diffeomorphisms satisfying the three conditions



Figure 1: The past $\mathcal{H}^-$ and future $\mathcal{H}^+$ horizons define the boundary of the Rindler wedge. Grey lines demonstrate lines of constant r (curved) and τ (straight). Long-wavelength perturbations of the hypersurface Σ_c are described by the equations of hydrodynamics.

i) The induced metric on the hypersurface Σ_c takes the form (10)

ii) The stress tensor on Σ_c describes a perfect fluid

iii) Diffeomorphisms return metrics stationary and homogeneous in (τ, x^i).

The allowed set is reduced to the following boost, shift and rescaling of x^μ. First, a constant boost β_i,

$$\sqrt{r_c}\tau \to \gamma(\sqrt{r_c}\tau - \beta_i x^i), \qquad x^i \to x^i - \gamma\beta^i\sqrt{r_c}\tau + (\gamma-1)\frac{\beta^i\beta_j}{\beta^2}x^j, \tag{13}$$

where $\gamma = (1-\beta^2)^{-1/2}$ and $\beta_i \equiv {r_c}^{-1/2}v_i$. Second, a shift in r and a rescaling of τ,

$$r \to r - r_h, \qquad \tau \to (1 - r_h/r_c)^{-1/2}\tau. \tag{14}$$

These yield the flat space metric in rather complicated coordinates,

$$\begin{aligned} ds^2 =& \frac{d\tau^2}{1-v^2/r_c}\left(v^2 - \frac{r-r_h}{1-r_h/r_c}\right) - \frac{2\gamma}{\sqrt{1-r_h/r_c}}d\tau dr - \frac{2\gamma v_i}{r_c\sqrt{1-r_h/r_c}}dx^i dr \\ &+ \frac{2v_i}{1-v^2/r_c}\left(\frac{r-r_c}{r_c-r_h}\right)dx^i d\tau + \left(\delta_{ij} - \frac{v_iv_j}{r_c^2(1-v^2/r_c)}\left(\frac{r-r_c}{1-r_h/r_c}\right)\right)dx^i dx^j. \end{aligned} \tag{15}$$

The Brown-York stress tensor on Σ_c (in units where $16\pi G = 1$) is given by

$$T_{ab} = 2(K\gamma_{ab} - K_{ab}), \tag{16}$$

where

$$K_{ab} = \frac{1}{2}(\mathcal{L}_n \gamma)_{ab}, \qquad K = K^a{}_a,$$

are the extrinsic curvature and its mean, and n^μ is the spacelike unit normal to the hypersurface.

By imposing that the Brown-York stress tensor on Σ_c gives that of the stress-energy tensor of a fluid we can identify the parameters of the metric (15) with the density, ρ, pressure, P and four-velocity u^a of a fluid, as follows:

$$\rho = 0, \qquad p = \frac{1}{\sqrt{r_c - r_h}}, \qquad u^a = \frac{1}{\sqrt{r_c - v^2}}(1, v_i). \tag{17}$$

The Hamiltonian constraint

$$R_{\mu\nu} n^\mu n^\nu = 0$$

on Σ_c yields a constraint on the Brown-York stress tensor

$$d T_{ab} T^{ab} = (T^a_a)^2.$$

When this constraint is applied to the equilibrium configurations described above, one finds the equation of state is $\rho = 0$ (as above), or $\rho = -2d(d-1)p$ which occurs for a fluid on the Taub geometry [14].

Promoting v_i and p to slowly varying functions of the the coordinates x^a, and regarding $v_i(\tau, x_j)$ and $p = r_c^{-1/2} + r_c^{-3/2} P(\tau, x_i)$ as small perturbations about equilibrium yields the metric

$$\begin{aligned} \mathrm{d}s^2 = &- r\mathrm{d}\tau^2 + 2\mathrm{d}\tau\mathrm{d}r + \mathrm{d}x^i\mathrm{d}x_i \\ &- 2\left(1 - \frac{r}{r_c}\right) v_i \mathrm{d}x^i\mathrm{d}\tau - 2\frac{v_i}{r_c}\mathrm{d}x^i\mathrm{d}r \\ &+ \left(1 - \frac{r}{r_c}\right)\left[(v^2 + 2P)\mathrm{d}\tau^2 + \frac{v_i v_j}{r_c}\mathrm{d}x^i\mathrm{d}x^j\right] + \left(\frac{v^2 + 2P}{r_c}\right)\mathrm{d}\tau\mathrm{d}r + \mathcal{O}(\epsilon^3) \end{aligned} \tag{18}$$

which satisfies the Einstein's equations to $\mathcal{O}(\epsilon^3)$ if v_i satisfies incompressibility, $\partial_i v^i = \mathcal{O}(\epsilon^3)$. Corrections appear in powers of ϵ^2, so this is the complete metric to second order.

The metric may now be built up order by order in the hydrodynamic scaling. Assume one has the metric at order ϵ^{n-1}, where the first non-vanishing component $\hat{R}^{(n)}_{\mu\nu}$ of the Ricci tensor appears at order n. By adding a correction term $g^{(n)}_{\mu\nu}$ to the metric at order n, resulting in a shift in the Ricci tensor $\delta R^{(n)}_{\mu\nu}$, and requiring

$$\hat{R}^{(n)}_{\mu\nu} + \delta R^{(n)}_{\mu\nu} = 0, \tag{19}$$

the vanishing of the Ricci tensor is guaranteed to order n. Recalling that, in the hydrodynamic scaling, derivatives scale thus,

$$\partial_r \sim \epsilon^0, \qquad \partial_i \sim \epsilon^1, \qquad \partial_\tau \sim \epsilon^2, \tag{20}$$

one sees that corrections $\delta R^{(n)}_{\mu\nu}$ at order n will appear only as r derivatives of $g^{(n)}_{\mu\nu}$. It is shown in [9] that, using the Bianchi identity and the Gauss-Codacci relations, integrability of the set of differential equations (19) defining $\delta R^{(n)}_{\mu\nu}$ in terms of $g^{(n)}_{\mu\nu}$ is given by imposing the momentum constraint, equivalent to the conservation of the stress tensor on Σ_c,

$$R_{a\mu}n^\mu = \nabla_a T^{ab}|^{(n)}_{\Sigma_c} = 0, \tag{21}$$

which is precisely the fluid equations of motion, to order n.

The perturbation scheme contains several degrees of freedom. The gauge freedom of the infinitesimal perturbations

$$g^{(n)}_{\mu\nu} \to g^{(n)}_{\mu\nu} + \partial_\mu \varphi^{(n)}_\nu + \partial_\nu \varphi^{(n)}_\mu$$

for some arbitrary vector $\varphi^{\mu(n)}(\tau, \vec{x}, r)$ at order ϵ^n, which may be fixed by demanding that $g_{r\mu}$ is that of the seed metric to all orders in ϵ. The x^a-dependent functions of integration from equation (19) may be fixed by imposing the boundary form (10) of the metric on Σ_c, and also requiring regularity of the metric at $r = 0$, which in this construction translates to the absence of logarithmic terms in r. Corrections to the bulk metric under these conditions then become

$$\begin{aligned}
g^{(n)}_{r\mu} =&0\\
g^{(n)}_{\tau\tau} =&(1 - r/r_c)F^{(n)}_\tau(\tau, \vec{x}) + \int_r^{r_c} \mathrm{d}r' \int_{r'}^{r_c} \mathrm{d}r''(\hat{R}^{(n)}_{ii} - r\hat{R}^{(n)}_{rr} - 2\hat{R}^{(n)}_{r\tau})\\
g^{(n)}_{\tau i} =&(1 - r/r_c)F^{(n)}_i(\tau, \vec{x}) - 2\int_r^{r_c} \mathrm{d}r' \int_{r'}^{r_c} \mathrm{d}r''\hat{R}^{(n)}_{ri}\\
g^{(n)}_{ij} =& - 2\int_r^{r_c} \mathrm{d}r'\frac{1}{r'} \int_0^{r'} \mathrm{d}r''\hat{R}^{(n)}_{ij},
\end{aligned} \tag{22}$$

where the $F^{(n)}_a(\tau, \vec{x})$ comprise of the remaining integration functions, and the final degree of freedom; field redefinitions of $\delta v^{(n)}_i$ and $\delta P^{(n)}$ at order ϵ^n. $F^{(n)}_i(\tau, \vec{x})$ is related to redefinitions of the fluid velocity and is fixed by the isotropic gauge condition $P^b_a T_{bc}u^c = 0$. $F^{(n)}_\tau(\tau, \vec{x})$ is related to redefinitions of the pressure and is fixed by defining the isotropic part of T_{ij} to be

$$T^{\text{isotropic}}_{ij} = \left(\frac{1}{\sqrt{r_c}} + \frac{P}{r_c^{3/2}}\right)\delta_{ij} \tag{23}$$

to all orders.

Applying the perturbation scheme to the seed metric yields to third order,

$$\begin{aligned}\mathrm{d}s^2 =& -r\mathrm{d}\tau^2 + 2\mathrm{d}\tau\mathrm{d}r + \mathrm{d}x^i\mathrm{d}x_i - 2\left(1-\frac{r}{r_c}\right)v_i\mathrm{d}x^i\mathrm{d}\tau - 2\frac{v_i}{r_c}\mathrm{d}x^i\mathrm{d}r \\ &+ \left(1-\frac{r}{r_c}\right)\left[(v^2+2P)\mathrm{d}\tau^2 + \frac{v_iv_j}{r_c}\mathrm{d}x^i\mathrm{d}x^j\right] + \left(\frac{v^2+2P}{r_c}\right)\mathrm{d}\tau\mathrm{d}r \\ &- \left[\frac{(r^2-r_c^2)}{r_c}\partial^2 v_i + \left(1-\frac{r}{r_c}\right)\left(\frac{v^2+2P}{r_c}\right)v_i\right]\mathrm{d}x^i\mathrm{d}\tau + \mathcal{O}(\epsilon^4)\end{aligned} \tag{24}$$

which satisfies the vacuum Einstein equations if

$$r_c^{3/2}\nabla^a T_{ai}|_{\Sigma_c} = \partial_\tau v_i - r_c\partial^2 v_i + \partial_i P + v^j\partial_j v_i = \mathcal{O}(\epsilon^5) \tag{25}$$

which are the Navier-Stokes equations with kinematical viscosity

$$\nu = r_c. \tag{26}$$

The corresponding corrections to the Navier-Stokes equations follow from conservation of the stress tensor on Σ_c. Vector and scalar quantities are odd and even orders respectively in the scaling ϵ. Accordingly, corrections to the scalar incompressibility equation appear at even orders, and to the vector Navier-Stokes equations at odd orders.

It is shown in [8] that the hydrodynamic limit is equivalently described on the Rindler wedge by taking the boundary to infinity $r_c \to \infty$, which effectively renders deviations from the horizon infinitesimal.

2.4 The Navier-Stokes fluid on a Schwarzschild sphere

In [10], Bredberg et al. construct the $(d+1)$-dimensional metric dual to the non-relativistic fluid on a d-dimensional hypersphere Σ_c^{S} (S denotes Schwarzschild) a distance $r_s > 0$ from the horizon of a Schwarzschild black hole. The Dirichlet boundary conditions used on the Rindler space are not consistent here. On Σ_c^{S}, one instead demands conformal flatness where the conformal factor is allowed to fluctuate, and constant mean curvature K defined with the unit tangent orthogonal to Σ_c^{S}. The perturbative expansion is parameterized by the inverse square of the curvature,

$$r_s = 1/4K^2,$$

in the limit $K \to \infty$. The metric is perturbed from the Schwarzschild metric

$$\mathrm{d}s^2_{\text{Schwarzschild}} = -\left[1-\left(\frac{2m}{\hat{r}}\right)^{(d-2)}\right]\mathrm{d}t^2 + \left[1-\left(\frac{2m}{\hat{r}}\right)^{(d-2)}\right]^{-1}\mathrm{d}\hat{r}^2 + \hat{r}^2\mathrm{d}\Omega^2_{d-1}. \tag{27}$$

where $\mathrm{d}\Omega_{d-1}^2$ is the metric on the unit $(d-1)$-sphere. In ingoing null coordinates (τ, r) defined by

$$\tau = \frac{(d-2)\alpha}{2m}\left(t + \int^{\hat{r}} \frac{\mathrm{d}\hat{r}'}{1-\left(\frac{2m}{\hat{r}'}\right)^{(d-2)}}\right), \quad r = \frac{2m}{\alpha(d-2)}(\hat{r}-2m), \tag{28}$$

the leading terms in the perturbed metric obeying the boundary conditions are

$$\begin{aligned}
\mathrm{d}s^2 = {} & -\frac{r}{r_s}\mathrm{d}\tau^2 + \frac{(d-1)(d-2)r(5+r)}{8m^2}\mathrm{d}\tau^2 + 2\mathrm{d}\tau\mathrm{d}r + 4m^2\mathrm{d}\Omega_{d-1}^2 \\
& + (1-r)[v^2\mathrm{d}\tau^2 - 2v_i\mathrm{d}\tau\mathrm{d}x^i] - 2rP\mathrm{d}\tau^2 + r_s\Big[(2(d-2)r + 8m^2P)\mathrm{d}\Omega_{d-1}^2 \\
& + (1-r)v_iv_j\mathrm{d}x^i\mathrm{d}x^j - (r^2-1)(\nabla^2 v_i + (d-2)R_{ji}v^j)\mathrm{d}\tau\mathrm{d}x^i - 2v_i\mathrm{d}r\mathrm{d}x^i \\
& + (v^2+2P)\mathrm{d}\tau\mathrm{d}r + 2(1-r)\hat{\phi}_i\mathrm{d}\tau\mathrm{d}x^i\Big] + r_s^2\Big(2\chi_i\mathrm{d}x^i\mathrm{d}r + \lambda_{ij}\mathrm{d}x^i\mathrm{d}x^j\Big) + \mathcal{O}(r_s^3)
\end{aligned} \tag{29}$$

for functions $(\hat{\phi}_i(x^a), \chi_i(r, x^a), \lambda_{ij}(r, x^a))$. The Ricci tensor on the $(d-1)$-sphere takes the form $R_{ij} = r_s^{-2}(d-2)g_{ij}$. The perturbed geometry satisfies Einstein's equations to $\mathcal{O}(r_s^0)$ if (v_i, P) satisfy the incompressible Navier-Stokes equations on the sphere with viscosity $\nu = 1$. In addition, the divergence of $\hat{\phi}_i(x^a)$ is fixed in terms of (v_i, P), and $(2\nabla^i\chi_i - \partial_\rho\lambda_i^i)$ is determined in terms of some function regular in r.

As Bredberg et al. point out, this spacetime points to a connection between famous conjectures on either side of the correspondence. The cosmic censorship conjecture states that singularities do not form on or outside event horizons in a spacetime arising from smooth initial data, while global existence for the Navier-Stokes similarly suggests that singularities do not form from smooth initial velocity fields. One can see that if a singularity were to form from smooth initial data, in the fluid velocity on Σ_c outside the horizon, this would simultaneously violate both conjectures.

3 Duality in the context of holography

The defining equations in general relativity are the Einstein field equations, and in the non-relativistic limit of hydrodynamics, the Navier-Stokes equations (9). Each is a set of non-linear partial differential equations describing the evolution of two fundamentally inextricable systems, in themselves exhibiting fantastically varied phenomenology. When approaching any complex physical system with a view to finding solutions, it is often advantageous to consider the symmetries, intensively studied in both of these systems since their conceptions. Beyond diffeomorphisms, the search in gravity has in general been somewhat limited [15, 36], however in the presence of a spacetime isometry, the symmetry group becomes remarkably large [35], particularly for vacuum spacetimes. For symmetries of the Navier-Stokes equations see [31], and with regards to the conformal group [12, 32]. In the light of the fluid/gravity correspondence, one may ask whether the symmetries of these systems are linked. In [37, 38, 39], they apply known symmetries of the Einstein equations to spacetimes with perfect fluid sources, constructing new spacetimes with the same equation of state, though not within a holographic framework. By drawing on the tools provided by these works and those in holography, we hope to develop a more general approach to the problem.

The bulk provides an additional valuable degree of freedom, where the boundary sets the scene for the fluid evolution on the induced geometry. Moreover, we are now free to exploit the symmetries of the more extensive yet simpler vacuum geometries. It is these symmetries which we intend to holographically project to the fluid. In particular, we are interested in dualities between solutions to the Navier-Stokes equations arising from dualities between solutions to the vacuum Einstein equations: dual metrics yield dual fluid configurations.

In this section, we discuss the work leading up to the spacetime Ehler's symmetry group of the vacuum field equations with zero cosmological constant. We continue in §4 to apply this to fluids on the boundary of the Rindler space, thus deriving solution generating transformations of the fluid velocity, pressure and viscosity (the latter defining the RG flow). We offer in §4.1.1 a selection of example transformations including RG flow for zero vorticity fluids (where one may relax this constraint), and $\mathbb{Z}_2$ transformations for fixed viscosity which we show in §4.2.2 in fact lie outside the spacetime Ehler's group.

3.1 Symmetry groups of the Einstein equations

Understanding the properties of the Einstein field equations has long been a subject of great theoretical interest, a sensible starting point being the inherent symmetries involved. To this end, Buchdahl [17] derived a form of duality in vacuum spacetime metrics, where an n-dimensional vacuum metric stationary with respect to a coordinate x^s:

$$g_{\mu\nu,s} = 0, \qquad g_{\mu s} = 0 \qquad \mu, \nu \in \{0, \ldots, n\}$$

generates a dual vacuum metric

$$h_{\mu\nu} = ((g_{ss})^{2/(n-3)} g_{ab}, (g_{ss})^{-1}), \qquad a, b \in \{\mu \neq s\}.$$

It is this solution generating property of spacetime isometries we wish to apply to solutions of the Einstein equations and holographically map to hydrodynamics.

We have, however, a considerably larger symmetry group at our disposal. The works of [16, 18, 19] develop the concept culminating in the generalized Ehlers symmetry group of the Einstein equations also for non-vacuum spacetimes. An extension exists [33, 34] to dualities between vacuum spacetimes and those with electromagnetic backgrounds described by the Einstein-Maxwell equations, of relevance for magnetohydrodynamics. We are concerned with the vacuum subgroup of the generalized Ehlers group, which includes the spacetime Ehlers group.

3.2 The spacetime Ehlers group

Let us begin by defining the Killing vector. Consider a manifold $\mathcal{M}$, endowed with a metric g. We define the set of Killing vectors $\xi \in T\mathcal{M}$ as the set of diffeomorphisms preserving the metric. This is equivalent to the statement that that Lie derivative of the metric along $\xi = \xi^\mu \partial_\mu$ vanishes:

$$(\mathcal{L}_\xi g)_{\mu\nu} = \xi^\rho g_{\mu\nu,\rho} + \xi^\rho{}_{,\mu}\, g_{\nu\rho} + \xi^\rho{}_{,\nu}\, g_{\mu\rho} = 0. \tag{30}$$

Consider a four-dimensional Lorentzian spacetime manifold with metric $g = (g_{\mu\nu})$ satisfying the vacuum Einstein equations, and admitting some Killing vector ξ. The twist potential

$$\omega_\mu = \sqrt{-\det(g)}\, \epsilon_{\mu\nu\sigma\rho} \xi^\nu \nabla^\sigma \xi^\rho \tag{31}$$

and Killing vector norm

$$\lambda = -\xi^\mu \xi_\mu \tag{32}$$

give the Ernst one-form

$$\sigma_\mu = \nabla_\mu \varsigma = \nabla_\mu \lambda - i\omega_\mu \tag{33}$$

for some scalar ς (exactness is guaranteed by vanishing Ricci tensor, see [40] p.164). Define a self-dual two form

$$\mathcal{F}_{\mu\nu} = (1 + i*)\nabla_\mu \xi_\nu\,, \tag{34}$$

where $*$ is the Hodge dual operator. Define a one form W satisfying

$$\nabla_{[\mu} W_{\nu]} = -2\gamma \Re[(\gamma\varsigma + i\delta)\mathcal{F}_{\mu\nu}] \tag{35a}$$

$$\xi^\alpha W_\alpha + 1 = (i\gamma\varsigma + \delta)(-i\gamma\bar{\varsigma} + \delta) = (\gamma\lambda)^2 + (\gamma\omega + \delta)^2 \tag{35b}$$

where a bar denotes complex conjugation, and γ and δ are non-simultaneously vanishing real constants, which as a pair fix the gauge of W. There exists a family of metrics, also with Killing vector ξ, which satisfy the vacuum Einstein equations, given by the map

$$g_{\mu\nu} \to h_{\mu\nu}(\xi, W, g) = \Omega^2 g_{\mu\nu} - 2\xi_{(\mu} W_{\nu)} - \frac{\lambda}{\Omega^2} W_\mu W_\nu \tag{36}$$

where $\Omega^2 \equiv \xi^\alpha W_\alpha + 1 \geq 1$ and the inequality holds over the whole geometry.

4 Solution-generating the Navier-Stokes fluid

It is clear that the dual metric $h(\xi, W, g)$ will in general take a different functional form from g. Consider the subset of transformations (36) preserving the functional form of $g(\phi)$ up to redefinitions $\phi \to \tilde{\phi}$ of some set ϕ of parameters. Vanishing of the Ricci tensor will yield a relation common to both metrics, in the parameters ϕ and their redefinitions.

In the case of the Rindler metric dual to the incompressible Navier-Stokes fluid, define these parameters by the fluid velocity v_i, pressure P, and boundary position r_c within the bulk. The *dual* metric is parametrised by a set of dual parameters $\tilde{v}_i$, $\tilde{P}$ and $\tilde{r}_c$, denoted by '$\sim$'. On satisfying the vacuum Einstein equations on $\tilde{\Sigma}_c$, now at $r = \tilde{r}_c$ in the dual geometry, the dual metric will yield the incompressible Navier-Stokes equations in the dual parameters

$$\partial_i \tilde{v}_i = 0 \tag{37a}$$

$$\partial_\tau \tilde{v}_i + \partial_i \tilde{P} + \tilde{v}_k \partial_k \tilde{v}_i - \tilde{r}_c \partial^2 \tilde{v}_i = 0. \tag{37b}$$

Vitally, if (v_i, P) satisfy the Navier-Stokes equations with viscosity $\nu = r_c$, then the dual velocity and pressure $(\tilde{v}_i, \tilde{P})$ represent a dual set of solutions for viscosity $\nu = \tilde{r}_c$. That is, a given subset of the spacetime Ehlers transformation acting on the fluid metric (24) obeying some Killing isometry corresponds to solution-generating transformations of the velocity and pressure, and RG flow parametrised by r_c, of an incompressible Navier-Stokes fluid.

At this point, we may generalise somewhat. The Rindler metric is just one fluid metric supporting flat background geometries on the boundary. We therefore wish instead to retain the common features of such metrics; the metric gauge $g_{\mu r}$, and the flat boundary metric which, by transforming to a suitable coordinate system, may always be expressed in the form (10). There is of course the additional freedom with vacuum metrics to introduce a constant conformal metric rescaling, varying under the duality, which we choose to neglect in our calculations. The equation we wish to solve is thus

$$g_{\mu\nu}(r_c, v_i, P) \to h_{\mu\nu}(\xi, W, g) = \tilde{g}_{\mu\nu} = g_{\mu\nu}(\tilde{r}_c, \tilde{v}_i, \tilde{P}). \tag{38}$$

where

$$\tilde{g}_{\tau r} = 1 + \frac{\tilde{v}(x^a)^2 + 2\tilde{P}(x^a)}{2\tilde{r}_c}, \qquad \tilde{g}_{ir} = -\frac{\tilde{v}_i(x^a)}{\tilde{r}_c}, \qquad \tilde{g}_{rr} = 0 \tag{39a}$$

$$\tilde{g}_{ab}|_{\tilde{r}_c} = \tilde{\gamma}_{ab}, \qquad \text{where} \qquad \tilde{\gamma}_{\tau\tau} = -\tilde{r}_c, \qquad \tilde{\gamma}_{ab} = \gamma_{ab} \qquad a \neq \tau \neq b. \tag{39b}$$

4.1 Transforming the fluid

We are provided in (39) with sufficient information to derive the fluid transformation via the form of the one-form W_μ. Preserving the vanishing of the $\tilde{g}_{rr} = g_{rr} = 0$ component of the metric we find, directly from (36), the two possibilities

$$W_r = -2\alpha\xi_r\Omega^2/\lambda \qquad \text{where} \quad \alpha = 0, 1. \tag{40}$$

One may obtain an expression for W_a by contraction of (36) with the boundary indices $(a, b, \ldots)$ of the Killing vector

$$W_a = \frac{\Omega^2\xi^r(g_{ar} + \xi_a\xi_r/\lambda) + \xi^b\tilde{g}_{ab}}{\lambda/\Omega^2 + (1-2\alpha)\xi^r\xi_r} - \frac{\Omega^2\xi_a}{\lambda}. \tag{41}$$

This expression is uniquely defined only at the dual boundary $\tilde{\Sigma}_c$, where we have defined the form of $\tilde{g}_{ab}$ and W_a becomes independent of the dual fluid velocity and pressure. These expressions diverge for null Killing vectors, where $\lambda = 0$. We cover this case shortly.

One can see that the definition of the fluid is given, to all orders in ϵ, by $g_{ar} = g_{ar}|_{r_c}$. Consequently, the fluid duality will be given by transformation of these components. Evaluation at $\tilde{r}_c$ is necessary in order to circumvent the ambiguity in the dual metric, and also provides explicit fluid transformations. We begin with Killing vectors null at the dual boundary, $\lambda|_{\tilde{r}_c} = 0$, where one finds from contraction of (36) with the Killing vector, which yields

$$\frac{\lambda}{\Omega^2}W_\mu = \xi^\nu(\tilde{g}_{\mu\nu} - g_{\mu\nu}), \tag{42}$$

the following transformation,

$$\tilde{g}_{ar} = g_{ar} + \left[\frac{\xi^b(g_{ab} - \tilde{\gamma}_{ab})}{\xi^r}\right]_{\tilde{r}_c}, \tag{43}$$

accompanied by the preservation of a null Killing vector, $\xi^\mu\xi^\nu\tilde{g}_{\mu\nu} = 0$.

For non-null Killing vectors we employ the relations

$$\xi^a(\tilde{g}_{ar} - (1-2\alpha)g_{ar}) = 0, \tag{44}$$

derived by comparing (40) and (42), and

$$\lambda/\Omega^2 = -\xi^\mu\xi^\nu\tilde{g}_{\mu\nu}, \tag{45}$$

found from contraction of (36) twice with the Killing vector. Inserting W_r (40) and W_a (41) into the Ehlers transformation (36) and employing (44) and (45), one finds

$$\tilde{g}_{ar} = \left[\frac{-\lambda g_{ar} + \xi_r((1-2\alpha)\xi^b\tilde{\gamma}_{ab} - \xi_a)}{\xi^c\xi^d\tilde{\gamma}_{cd} + (1-2\alpha)\xi^r\xi_r}\right]_{\tilde{r}_c}. \tag{46}$$

4.1.1 Energy scaling invariance from a bulk isometry

We begin with an example a (null) Killing vector into the bulk, $\xi = \xi^r(x^\mu)\partial_r$. Integrability of the Killing equations requires

$$2v_j\partial_{[i}v_{j]} = -r_c\partial^2 v_i \tag{47}$$

along with vanishing vorticity, which with incompressibility implies (47). Transformation (43) gives

$$\tilde{v}_i = \frac{\tilde{r}_c}{r_c}v_i, \quad \tilde{P} = \frac{\tilde{r}_c}{r_c}P + \frac{\tilde{r}_c}{r_c}\left(1 - \frac{\tilde{r}_c}{r_c}\right)\frac{v^2}{2} \tag{48}$$

It is trivial to show that the pair $(\tilde{v}_i, \tilde{P})$ satisfy the incompressible Navier-Stokes equations if (v_i, P) do so for velocities satisfying (47) alone - vanishing vorticity imposes unnecessary constraint and removes the dissipative term from the fluid equations.

It is interesting to consider the problems of existence, uniqueness and regularity of the Navier-Stokes in this case. The divergence of (47) yields a vanishing mean square vorticity which ensures the class of solutions (v_i, P) generated by (48) are regular. With respect to existence, the kinetic energy scales by a factor $\tilde{r}_c/r_c$ and thus is bounded if there exists any solution satisfying (47) where the energy is finite.

We note that the twist vanishes, and there exist values of γ/δ for which $\Omega^2 = 0$ and the dual metric exhibits curvature singularities. We do not know how, if at all, this translates to the dual fluid.

4.1.2 The timelike Killing vector

One might expect from the timelike Killing vector $\xi = \partial_\tau$ a duality of the form

$$\tilde{v}_i = -v_i, \qquad \tilde{P} = P, \qquad \tilde{r}_c = -r_c \tag{49}$$

enacting time-reversal of the fluid but this is not the case. This is explained by noting that time-reversal is enacted by redefining the viscosity by $\nu = \pm r_c$ [8] rather than by changing r_c itself. This is because sending $r_c \to -r_c$ brings the fluid outside the causal region of the spacetime.

4.2 Fixed viscosity dualities

We turn to fixed boundary (viscosity) transformations, where $\tilde{r}_c = r_c$. For Killing vectors null at the dual boundary, or if $\alpha = 0$, one recovers the identity. For non-null Killing vectors with $\alpha = 1$, one finds

$$\tilde{g}_{ar} = \left[\frac{g_{ar} - 2\xi_r \xi^b \gamma_{ab} / \xi^e \xi^f \gamma_{ef}}{1 - \xi^r \xi_r / \xi^c \xi^d \gamma_{cd}} \right]_{r_c}, \tag{50}$$

which constitutes a $\mathbb{Z}_2$ group.

4.2.1 Spacelike Killing vectors

Consider a generic space-like Killing vector $\xi = \xi^k \partial_k$. Under (50), the pressure is preserved, while the velocity transforms as

$$\tilde{v}_i = v_i - 2\xi_i \frac{\sum_k \xi^k v_k}{\sum_j (\xi^j)^2}, \tag{51}$$

which is a reflection in the hyperplane normal to the Killing vector and containing the point at which the velocity is defined.

Translational isometry

Consider $\xi = c_k \partial_k$ where c_k are normalised to $\sum_k c_k^2 = 1$, and the corresponding isometries are $c_k \partial_k v_i = c_k \partial_k P = 0$. The dual fields are

$$\tilde{v}_i = v_i - 2c_i c_k v_k \qquad \tilde{P} = P. \tag{52}$$

The incompressibility condition

$$\partial_i \tilde{v}_i = \partial_i v_i - 2c_i \partial_i c_k v_k = 0, \tag{53}$$

and Navier-Stokes equations

$$\begin{aligned} \partial_\tau \tilde{v}_i + \partial_i \tilde{P} + \tilde{v}_k \partial_k \tilde{v}_i - r_c \partial^2 \tilde{v}_i =& (\delta_{ik} - 2c_i c_k)(\partial_\tau v_k + \partial_k P + v^j \partial_j v_k - r_c \partial^2 v_k) \\ &+ 2c_i c_k \partial_k P - 2c_j v_j c_k \partial_k (v_i - 2c_i c_l v_l) = 0, \end{aligned} \tag{54}$$

are satisfied by the incompressible Navier-Stokes equations in the original fluid parameters along with the isometries. This is valid for fluids of arbitrary dimension.

Rotational isometry

Consider a Killing vector $\xi = -x_2 \partial_1 + x_1 \partial_2$ corresponding to a rotational isometry in a d-dimensional fluid. In polar coordinates

$$x_1 = \rho \cos\theta, \qquad x_2 = \rho \sin\theta \qquad x_{k'} = x_{k'}, \tag{55}$$

where primed ($'$) indices run from 3 to $(d-1)$, and the Killing vector becomes $\xi = \partial_\theta$, the isometries are

$$\partial_\theta v_1 = -v_2 \qquad \partial_\theta v_2 = v_1 \qquad \partial_\theta v_{k'} = 0 \qquad \partial_\theta P = 0. \tag{56}$$

Solutions satisfy

$$v_1 = x_1 \mu(\tau, x_{k'}, \rho) - x_2 \eta(\tau, x_{k'}, \rho) \qquad v_2 = x_2 \mu(\tau, x_{k'}, \rho) + x_1 \eta(\tau, x_{k'}, \rho) \tag{57}$$

with dual velocities

$$\tilde{v}_1 = x_1 \mu(\tau, x_{k'}, \rho) + x_2 \eta(\tau, x_{k'}, \rho) \qquad \tilde{v}_2 = x_2 \mu(\tau, x_{k'}, \rho) - x_1 \eta(\tau, x_{k'}, \rho), \tag{58}$$

that is, the transformation sends $\eta \to -\eta$ (equivalently $\theta \to -\theta$). The incompressible Navier-Stokes equations for the original fluid may be expressed as

$$\begin{aligned}
0 &= (2 + \rho\partial_\rho)\mu + \partial_{k'} v_{k'} && \text{(59a)}\\
0 &= \partial_\tau \mu + \rho^{-1}\partial_\rho P + \mu^2 - \eta^2 + \mu\rho\partial_\rho \mu + v_{k'}\partial_{k'}\mu && \text{(59b)}\\
&\qquad - r_c(3\rho^{-1}\partial_\rho \mu + \partial_\rho^2 \mu + \partial^{k'}\partial_{k'}\mu) &&\\
0 &= \partial_\tau \eta + 2\mu\eta + \mu\rho\partial_\rho \eta + v_{k'}\partial_{k'}\eta - r_c(3\rho^{-1}\partial_\rho \eta + \partial_\rho^2 \eta + \partial^{k'}\partial_{k'}\eta) && \text{(59c)}\\
0 &= \partial_\tau v_{k'} + \partial_{k'} P + \mu\rho\partial_\rho v_{k'} + v_{j'}\partial_{j'} v_{k'} - r_c \rho^{-1}\partial_\rho(\rho\partial_\rho v_{k'}) && \text{(59d)}
\end{aligned}$$

It is clear from the parity of these equations in η that if there exists a fluid solution defined in terms of a pair (μ, η) by (57), then there also exists a solution parameterized by the pair $(\mu, -\eta)$. That is, the dual fluid satisfies the incompressible Navier-Stokes equations. Again, this is valid for fluids of arbitrary dimension.

We provide an example with the three-dimensional fluid solution

$$\begin{aligned}
v_1 &= A\left(x_1 - x_2 e^{-2A(\tau-\tau_0)}\right) \qquad v_2 = A\left(x_2 + x_1 e^{-2A(\tau-\tau_0)}\right) && \text{(60a)}\\
v_3 &= B\exp\left(4A(\tau-\tau_0) + \frac{A\rho^2}{2r_c}\right) - e^{2A\tau}\int \mathrm{d}\tau'\, q(\tau') e^{-2A\tau'} - 2Ax_3 && \text{(60b)}\\
P &= \frac{1}{2}A^2\rho^2(e^{-4A(\tau-\tau_0)} - 1) + q(\tau)x_3 - 2A^2 x_3^2 && \text{(60c)}
\end{aligned}$$

which satisfies the isometries (56). Here, A, B and τ_0 are arbitrary non-vanishing constants and $q(\tau)$ is an arbitrary function of time. The duality is equivalent to sending $\tau_0 \to \tau_0 + i\pi/2A$.

4.2.2 Group structure for fixed boundary

Consider the fixed boundary transformations in the case $\xi^r = 0$. The constraints (35a) on $\partial_{[r}W_{a]}$ are dependent on radial derivatives of the dual metric, which we have not defined, and so remain arbitrary. The RHS of (35a) is closed [16], so it remains to constrain $\partial_{[a}W_{b]}|_{r_c}$, which vanishes. Now, we may write the Killing vector norm in the form

$$-\lambda = \gamma_{ab}\xi^a\xi^b + (r_c - r)\{\xi^\tau(g_{\tau r} + \mathcal{O}(\epsilon^4)) + \xi^k(g_{kr} + \mathcal{O}(\epsilon^3))\}^2, \tag{61}$$

which by (44) is preserved, up to fourth order terms, under the duality. Equation (45) then yields $\Omega^2 = 1+\mathcal{O}(\epsilon^4)$, and the constraints are satisfied for $\gamma = 0$, $\delta = \pm 1$.

One begins in four dimensions [16] with the spacetime Ehlers group isomorphic to $SL(2,\mathbb{R})$, parametrised in terms of the Möbius map of the Ernst scalar $\varsigma \to (\delta'\varsigma + i\gamma')/(i\gamma\varsigma + \delta)$ by the real constants $(\gamma, \delta, \gamma', \delta')$ satisfying $\gamma'\gamma + \delta'\delta = 1$. For fixed boundary, $\gamma = 0$ and $\delta = \pm 1$ in the Möbius map result in a constant imaginary shift in the Ernst scalar which can be absorbed as an arbitrary integration constant of the twist. In this way, the metric transformation is decoupled from the $SL(2,\mathbb{R})$ action.

Transformation (36) forms a group acting on metrics in arbitrary spacetime dimension [16], which is not in general vacuum-to-vacuum, but contains the four dimensional spacetime Ehlers group as a particular case. The fixed boundary fluid transformations arise from vacuum-to-vacuum maps within the generalized Ehler's group, but outside the spacetime Ehler's group. For those examples in §4.2.1, explicit calculation of the action of (51) on the Ernst scalar in four spacetime dimensions yields $\varsigma \to \bar{\varsigma}(1 + \mathcal{O}(\epsilon^4))$ (provided the arbitrary integration constant of the twist vanishes), which provides some hint as to the form of this spacetime transformation. Note, in the presence of non-vanishing twist, this complex conjugation is not found in the Möbius map for any values of the group parameters.

5 Discussion

We have demonstrated how solution-generating transformations of the vacuum Einstein equations in the presence of a Killing vector may be applied to spacetimes holographically dual to hydrodynamics. Our focus has been on the incompressible Navier-Stokes fluid dual to vacuum Rindler spacetime, where we have uncovered a selection of dualities: a linear energy scaling symmetry for solutions with vanishing vorticity (this constraint may be relaxed to (48)), deriving from RG flow of the fluid hypersurface through the bulk, and a $\mathbb{Z}_2$ group of transformations for fixed viscosity (boundary) with explicit examples of reflection-like symmetry in translational and rotational fluid isometries. These transformations may not be remarkable from the perspective of hydrodynamics but it shows how part of the Ehler's transformations can survive holography and give rise to transformations in the fluid dual. One can then be inspired to try the Harrison transformation which is a solution generating transformation in Einstein-Maxwell theory to give new transformations in magnetohydrodynamics. A holographic relation of the sort described here has been constructed for magnetohydrodynamics in [41, 42]. The Harrison transformation in the bulk may then lead to nontrivial transformations in magnetohydrodynamics transforming between fluid velocity and magnetic potentials; this is the subject of current work. One can also study the dimensional dependence of these solution generating transformations. In recent work [43], the difference in scaling for turbulence between three and four dimensions was studied holographically with a large difference in qualitative behaviour. One could also examine backgrounds associated with nontrivial chemical potential in the dual; for example rotating or charged black hole backgrounds.

Essentially, in this paper we wish to open up the use of gravitational solution generating symmetries in holography. It is encouraging that this did not immediately fail and one could preserve the fluid metric ansatz with some residual transformations surviving, yet it is intriguing that these transformations did not give anything particularly new. The results for magnetohydrodynamics may prove more significant.

Acknowledgements

We would like to thank Cynthia Keeler and Andrew Strominger for their illuminating and insightful thoughts and comments on the fluid gravity correspondence and Shiraz Minwalla whose lectures on hydrodynamics at the Isaac Newton Institute programme on M-theory were an inspiration. This work is partially supported by STFC consolidated grant ST/J000469/1.

References

[1] T. Damour, *Quelques propriétés mécaniques, électromagnétiques, thermodynamiques et quantiques des trous noirs*, Thèse de Doctorat d'Etat, Université Pierre et Marie Curie, Paris VI, 1979.

[2] K. S. Thorne, R. H. Price, D. A Macdonald, *Black Holes: The Membrane Paradigm*, Yale University Press, 1986

[3] R. Baier, P. Romatschke, D. T. Son, A. O. Starinets and M. A. Stephanov, *Relativistic viscous hydrodynamics, conformal invariance, and holography*, JHEP **0804** (2008) 100 [arXiv:0712.2451 [hep-th]].

[4] P. Kovtun, D. T. Son and A. O. Starinets, *Viscosity in strongly interacting quantum field theories from black hole physics*, Phys. Rev. Lett. **94** (2005) 111601 [hep-th/0405231].

[5] G. Policastro, D. T. Son and A. O. Starinets, *The Shear viscosity of strongly coupled N=4 supersymmetric Yang-Mills plasma*, Phys. Rev. Lett. **87** (2001) 081601 [hep-th/0104066].

[6] S. Bhattacharyya, V. EHubeny, S. Minwalla and M. Rangamani, *Nonlinear Fluid Dynamics from Gravity*, JHEP **0802** (2008) 045 [arXiv:0712.2456 [hep-th]].

[7] I. Bredberg, C. Keeler, V. Lysov and A. Strominger, *Wilsonian Approach to Fluid/Gravity Duality*, JHEP **1103** (2011) 141 [arXiv:1006.1902 [hep-th]].

[8] I. Bredberg, C. Keeler, V. Lysov and A. Strominger, *From Navier-Stokes To Einstein*, JHEP **1207** (2012) 146 [arXiv:1101.2451 [hep-th]].

[9] G. Compere, P. McFadden, K. Skenderis and M. Taylor, *The Holographic fluid dual to vacuum Einstein gravity*, JHEP **1107** (2011) 050 [arXiv:1103.3022 [hep-th]].

[10] I. Bredberg and A. Strominger, *Black Holes as Incompressible Fluids on the Sphere*, JHEP **1205** (2012) 043 [arXiv:1106.3084 [hep-th]].

[11] R. -G. Cai, L. Li and Y. -L. Zhang, *Non-Relativistic Fluid Dual to Asymptotically AdS Gravity at Finite Cutoff Surface*, JHEP **1107** (2011) 027 [arXiv:1104.3281 [hep-th]].

[12] S. Bhattacharyya, S. Minwalla and S. R. Wadia, *The Incompressible Non-Relativistic Navier-Stokes Equation from Gravity*, JHEP **0908** (2009) 059 [arXiv:0810.1545 [hep-th]].

[13] G. Compere, P. McFadden, K. Skenderis and M. Taylor, *The relativistic fluid dual to vacuum Einstein gravity*, JHEP **1203** (2012) 076 [arXiv:1201.2678 [hep-th]].

[14] C. Eling, A. Meyer and Y. Oz, *The Relativistic Rindler Hydrodynamics*, JHEP **1205** (2012) 116 [arXiv:1201.2705 [hep-th]].

[15] C. G. Torre and I. M. Anderson, *Symmetries of the Einstein equations*, Phys. Rev. Lett. **70** (1993) 3525 [gr-qc/9302033].

[16] M. Mars, *Space-time Ehlers group: Transformation law for the Weyl tensor*, Class. Quant. Grav. **18** (2001) 719 [gr-qc/0101020].

[17] H. A. Buchdahl, *Reciprocal Static Metrics and Scalar Fields in the General Theory of Relativity*, Phys. Rev. **115** (1959) 1325. `http://prola.aps.org/abstract/PR/v115/i5/p1325_1`.

[18] J. Ehlers, *Konstruktionen und Charakterisierung von Losungen der Einsteinschen Gravitationsfeldgleichungen.*

[19] R. P. Geroch, *A Method for generating solutions of Einstein's equations*, J. Math. Phys. **12** (1971) 918. doi:10.1063/1.1665681.

[20] T. Padmanabhan, *Entropy density of spacetime and the Navier-Stokes fluid dynamics of null surfaces*, Phys. Rev. D **83** (2011) 044048 [arXiv:1012.0119 [gr-qc]].

[21] Y. Matsuo, M. Natsuume, M. Ohta and T. Okamura, *The Incompressible Rindler fluid versus the Schwarzschild-AdS fluid*, arXiv:1206.6924 [hep-th].

[22] S. Chapman, Y. Neiman and Y. Oz, *Fluid/Gravity Correspondence, Local Wald Entropy Current and Gravitational Anomaly*, JHEP **1207** (2012) 128 [arXiv:1202.2469 [hep-th]].

[23] T. -Z. Huang, Y. Ling, W. -J. Pan, Y. Tian and X. -N. Wu, *Fluid/gravity duality with Petrov-like boundary condition in a spacetime with a cosmological constant*, Phys. Rev. D **85** (2012) 123531 [arXiv:1111.1576 [hep-th]].

[24] F. G. Rodrigues, W. A. Rodrigues, Jr and R. da Rocha, *The Maxwell and Navier-Stokes that Follow from Einstein Equation in a Spacetime Containing a Killing Vector Field*, AIP Conf. Proc. **1483** (2012) 277 [arXiv:1109.5274 [math-ph]].

[25] R. Nakayama, *The Holographic Fluid on the Sphere Dual to the Schwarzschild Black Hole*, arXiv:1109.1185 [hep-th].

[26] T. -Z. Huang, Y. Ling, W. -J. Pan, Y. Tian and X. -N. Wu, *From Petrov-Einstein to Navier-Stokes in Spatially Curved Spacetime*, JHEP **1110** (2011) 079 [arXiv:1107.1464 [gr-qc]].

[27] D. Brattan, J. Camps, R. Loganayagam and M. Rangamani, *CFT dual of the AdS Dirichlet problem : Fluid/Gravity on cut-off surfaces*, JHEP **1112** (2011) 090 [arXiv:1106.2577 [hep-th]].

[28] S. Kuperstein and A. Mukhopadhyay, *The unconditional RG flow of the relativistic holographic fluid*, JHEP **1111** (2011) 130 [arXiv:1105.4530 [hep-th]].

[29] L. D. Landau, E. M. Lifschitz, *Fluid Mechanics, Course of Theoretical Physics*, Oxford: Pergamon Press, 1959.

[30] M. Rangamani, *Gravity and Hydrodynamics: Lectures on the fluid-gravity correspondence*, Class. Quant. Grav. **26** (2009) 224003 [arXiv:0905.4352 [hep-th]].

[31] V. N. Gusyatnikova, V. A. Yumaguzhin, *Symmetries and conservation laws of navier-stokes equations*, Acta Applicandae Mathematicae **15** (January, 1989) 65-81.

[32] P. A. Horvathy and P. -M. Zhang, *Non-relativistic conformal symmetries in fluid mechanics*, Eur. Phys. J. C **65** (2010) 607 [arXiv:0906.3594 [physics.flu-dyn]].

[33] M. L. Woolley *On the role of the Killing tensor in the Einstein-Maxwell theory* Volume 33, Number 2 (1973), 135-144, doi: 10.1007/BF01645625.

[34] B. K. Harrison, *New Solutions of the Einstein-Maxwell Equations from Old* J. Math. Phys. 9, 1744 (1968); doi: 10.1063/1.1664508.

[35] D. Maison in *Duality and Hidden Symmetries in Gravitational Theories*, in *Einstein's Field Equations and Their Physical Implications*, B. G. Schmidt (ed.), Springer-Verlag, 2000.

[36] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, E. Herlt, *Exact Solutions of Einstein's Field Equations*, Second Edition. 732 p., Cambridge University Press, Cambridge, 2003. GBP80.00, ISBN 0521461367.

[37] H. Stephani, *Symmetries of Einstein's field equations with a perfect fluid source as examples of Li-Backlünd symmetries*, J. Math. Phys. 29, 1650 (1988); doi: 10.1063/1.527913.

[38] D. Garfinkle, E. N. Glass and J. P. Krisch, *Solution generating with perfect fluids*, Gen. Rel. Grav. **29** (1997) 467 [gr-qc/9611052].

[39] P. Boonserm, M. Visser and S. Weinfurtner, *Solution generating theorems for perfect fluid spheres*, J. Phys. Conf. Ser. **68** (2007) 012055 [gr-qc/0609088].

[40] R. M. Wald, *General Relativity*, Chicago, Usa: Univ. Pr. (1984) 491p.

[41] C. -Y. Zhang, Y. Ling, C. Niu, Y. Tian and X. -N. Wu, *Magnetohydrodynamics from gravity* Phys. Rev. D **86** (2012) 084043 arXiv:1204.0959 [hep-th].

[42] C. Niu, Y. Tian, X. -N. Wu and Y. Ling, *Incompressible Navier-Stokes Equation from Einstein-Maxwell and Gauss-Bonnet-Maxwell Theories*, Phys. Lett. B **711** (2012) 411 [arXiv:1107.1430 [hep-th]].

[43] F. Carrasco, L. Lehner, R. C. Myers, O. Reula and A. Singh, *Turbulent flows for relativistic conformal fluids in 2+1 dimensions* arXiv:1210.6702 [hep-th].